

Agricultural Research Institute, Pusa

Sampling for Rice Yield in Bihar and Orissa

BY

J. A. HUBBACK, I.C.S.



CALCUTTA : GOVERNMENT OF INDIA
CENTRAL PUBLICATION BRANCH

1927

**Government of India Publications are obtainable from the
Government of India Central Publication Branch,
3, Government Place, West, Calcutta, and from
the following Agents :—**

EUROPE.

OFFICE OF THE HIGH COMMISSIONER FOR INDIA,
42, GROSVENOR GARDENS, LONDON, S.W. 1.
And at all Booksellers.

INDIA AND CEYLON.

Provincial Book Depôts:

MADRAS :—Office of the Superintendent, Government Press, Mount Road, Madras.
BOMBAY :—Superintendent, Government Book Depot, Town Hall, Bombay.
SIND :—Library attached to the Office of the Commissioner in Sind, Karachi.
JENGAH :—Office of the Bengal Secretariat Book Depot, Writers' Buildings, Room No. 1, Ground Floor, Calcutta.
UNITED PROVINCES OF AGRA AND OUDH :—Office of the Superintendent of Government Press, United Provinces of Agra and Oudh, Allahabad.
PUNJAB :—Office of the Superintendent, Government Printing, Punjab, Lahore.
TERMA :—Office of the Superintendent, Government Printing, Burma, Rangoon.
CENTRAL PROVINCES AND BEHAR :—Office of the Central Provinces Secretariat, Nagpur.
ASSAM :—Office of the Superintendent, Assam Secretariat Press, Shillong.
BIHAR AND ORISSA :—Office of the Superintendent, Government Printing, Bihar and Orissa, P. O., Gulzarbagh, Patna.
COORG :—Office of the Chief Commissioner of Coorg, Bangalore.
NORTH-WEST FRONTIER PROVINCE :—Office of the Manager, Government Printing and Stationery, Peshawar.

Thacker, Spink & Co., Calcutta and Simla.
W. Newman & Co., Ltd., Calcutta.
R. Cambray & Co., Calcutta.
S. K. Lahiri & Co., Calcutta.
The Indian School Supply Depot, 309, Bow Bazar Street, Calcutta, and 220, Nawab-pur, Dacca.
Butterworth & Co. (India), Ltd., Calcutta.
Rai M. C. Sarcar Bahadur & Sons, 90-2A, Harrison Road, Calcutta.
The Vedion Library, 17, Park Street, Calcutta.
Standard Literature Company, Limited, Calcutta.
Association Press, Calcutta.
Chukerverty, Chatterjee & Co., Ltd., 13, College Square, Calcutta.
The Book Company, Calcutta.
James Murray & Co., 12 Government Place, Calcutta (for Meteorological Publications only).
Ray Choudhury & Co., 65-5, Asutosh Mukherjee Road, Calcutta.
The Oriental Publishing House, 34, Cornwallis Street, Calcutta.
D. C. Dasak 184, Proprietor, Albert Library, Dacca.
Hightbotham, Madras.
V. Kalyanaswami Iyer & Co., Madras.
P. R. Rama Iyer & Co., Madras.
Rohouse and Sons, Madras.
G. A. Robinson & Co., Publishers, George Town, Madras.
The Modern Stores, Salem, Madras.
Theosophical Publishing House, Adyar, Madras.
Bright & Co., Trivandrum.
The Booklover's Resort, Takad, Trivandrum, South India.
V. S. Swaminathan, Bookseller, West Tower Street, Madras.
E. M. Gopalakrishna Kone, Podumandapam, Madras.
Vijapur & Co., Vizagapatam.
Thacker & Co., Ltd., Bombay.
D. B. Anagorewalla, Sons & Co., Bombay.
Sunder Pandurang, Bombay.
Ran Chandra Govind & Sons, Kalbadevi, Road, Bombay.
N. M. Tripathi & Co., Booksellers, Princes Street, Kalbadevi Road, Bombay.
Proprietor, New Kitabhana, Poona.
The Manager, Oriental Book Supplying Agency, 15, Shikharwar, Poona City.
R. S. Gondhalekar's Book Depot, Publisher and Bookseller, Budhwar Chawk, Poona City.
Managing Director, Co-operative Bookstall, Booksellers and Publishers, Poona City.
Rama Krishna Bros., Opposite Viarimbag, Poona City.
Karanandas Narandas & Sons, Surat.
Mahajadas & Sons, Booksellers and Publishers, Bhaga Talao, Surat.
Mrs. Radhnabi Atmaram Sagoon, Kalbadevi Road, Bombay.

A. H. Wheeler & Co., Allahabad, Calcutta and Bombay.
R. B. Unadikar & Co., the Bharat Book Depot, Dharwar.
The Standard Bookstall, Karachi, Quetta, Delhi, Murree and Rawalpindi.
The Karachi Book Depot, Elphinstone Street, Camp, Karachi.
The Standard Bookstall, Quetta.
U. P. Mahindra & Co., Quetta.
J. Ray & Sons, 43, R. & L. Edwardes Road, Rawalpindi and Murree.
The Standard Book Depot, Lahore, Lucknow, Nainital, Mussoorie Dalhousie, Ambala Cantonment and Delhi.
N. B. Mathur, Superintendent, Nazir Khami Hind Press, Allahabad.
The North India Christian Tract and Book Society, 18, Clive Road, Allahabad.
Ram Dayal Agarwala, 184, Katra, Allahabad.
Manager, Newal Kishore Press, Lucknow.
The Upper India Publishing House, Ltd., 41, Amnabad Park, Lucknow.
Munshi Soela Ram, Managing Proprietor, Indian Army Book Depot, Juhl, Cawnpore.
Rai Sahib M. Gulab Singh & Sons, Muddi-Am Press, Lahore and Allahabad.
Ramu Kriehna & Sons, Booksellers, Anarkali, Lahore.
Puri Brothers, Booksellers and Publishers, Kacheri Road, Lahore.
The Tilak School Bookshop, Lahore.
The Standard Bookstall, Lahore.
Manager of the Imperial Book Depot, 63, Chandney Chawk Street, Delhi.
Oxford Book and Stationery Company, Delhi.
Superintendent, American Baptist Mission Press, Rangoon.
Proprietor, Rangoon Times Press, Rangoon.
The Modern Publishing House, Ltd., 50, Phayre Street, Rangoon.
The International Buddhist Book Depot, Post Box No. 971, Rangoon.
Burma Book Club, Ltd., Rangoon.
Manager, the "Hitasava," Nagpur.
Bhisey Brothers, Booksellers and Stationers, Sitabaldi, Nagpur.
S. C. Talukdar, Proprietor, Students & Co., Cooch Behar.
Times of Ceylon Co., Ltd., Colombo.
The Manager, Ceylon Observer, Colombo.
The Manager, The Indian Book Shop, Benares City.
The Srivilliputtur Co-operative Trading Union, Ltd., Srivilliputtur (Satur S. I. R.).
Banwari Lal, Esq., Fakarya Street, Pilibhit, United Provinces.
The Manager, Educational Book Depot, Jubbulpore.
Raghunath Prasad & Sons, Patna City.
Dandekar, Brothers, Indore City.
The Hyderabad Book Depot, Chaderghat, Hyderabad (Deccan).

Sampling for Rice* Yield in Bihar and Orissa.

(Received for publication on 10th November 1926.)

The need for reasonably accurate statistics of the human population of a country has long been recognized throughout the civilized world. The idea that statistics of the production which supports that population are of well-nigh equal importance is rapidly gaining general recognition. It is, therefore, perhaps superfluous to argue in its favour, but it may be useful to indicate what objects a well-founded estimate of the yield of rice can fulfil.

In the first place, since rice occupies a little more than half of the cultivated area in the province of Bihar and Orissa, employs 4-5th of the total population for six months of the year and feeds the greater part of it throughout the year, its supreme importance in any survey of production can scarcely be denied. It probably provides at least one-half of the rural contribution to the "provincial dividend", and that contribution is still vastly more important than that of the urban and industrial workers. Hence if a survey of production is desirable, good statistics of rice yield are essential.

Apart from the uses to which those who direct or influence the policy of the State may put the statistics of production, there is no doubt that traders want information of crops as they come forward. Good information enables them to operate with more confidence and not only saves them from alternating between prosperity and financial disaster, but greatly benefits the public by stabilizing prices. For this purpose it is not enough to know what is the production of the whole province, but knowledge of the conditions of the districts or even smaller units is very valuable. Lastly, reasonably accurate knowledge of the real normal yields of important crops even for fairly large tracts would give guidance to courts who are required to estimate the value of produce for the decision of rent suits, for commuting produce rent or for settling fair cash rents.

* I have used the word "rice" throughout as the ordinary description of the crop. As a matter of fact the samples taken are, strictly, "paddy" i.e. unhusked rice.

Hitherto lack of such knowledge has frequently led to considerable injustice.

The statistics of area under the rice crop are claimed to be fairly accurate. They are founded ultimately on the figures obtained from detailed measurement in a particular year on the ground. No doubt new land has been brought under rice in some places, the year of survey may have been a particularly favourable or unfavourable one, or to a limited extent other crops may have replaced rice. Allowance is made for these factors year by year, and perhaps there is no very large error. The question is examined further in the latter part of this bulletin. But in the figure for yield there is certainly room for bad mistakes. This has been fixed under the advice of the Director of Agriculture at a definite figure as normal, and is varied year by year by applying for each district a percentage, indicating how far the crop of the year falls below or rises above the normal. Now in the first place it is exceedingly probable that the normal rice yield differs materially between districts. But, apart from that, there is no machinery for fixing with any reasonable accuracy the percentage of the year. All that is done at present is for the local police officers to make a guess, at which in succession the Subdivisional Officer, the District Officer, and the Director of Agriculture guess again. When it is considered that the percentage depends ultimately on the effect of the weather on very various soils, cultivated with varying degrees of skill and enterprise, planted with different kinds of rice, protected by irrigation works of greater or less efficiency or completely unprotected, liable to or immune from crop pests, and finally harvested over a period of nearly three months, it becomes apparent that the guessing ability of the officers concerned has to be remarkable. Unfortunately, too, not one of them has the least chance of finding out whether his guess was fairly right or wildly wrong. Hence the existing statistics of rice production are, I believe, the result of applying to a fairly accurate figure of area, an arbitrary standard of normal yield, and a pure guess of the condition of the year. I am convinced that by an expenditure of a sum of money, that is not excessive in view of the importance of the matter, the last two and perhaps the first could be replaced by a scientific estimate, in which the possibilities of error can be gauged. Incidentally the scheme proposed, though it would not completely solve the problems of the courts to which reference has been made, would go a great way to do so.

I return to the point of the normal yield. This is based on a number of crop-cutting experiments made at different times in different places,

and it is to the defects of these experiments, with the consequent uncertainty of the conclusions based on them, that I now wish to draw attention.

The commonest method of estimating yield, apart from pure guessing from the appearance of the field at harvest time, is that described in the standard account of the *danabandi* (appraisement) system of rent payment. It consists of the tenant cutting a small area where the crop is thinnest and the landlord where the crop is heaviest, mixing the two together and taking the result as a fair sample. To secure accurate results this presupposes that the true mean lies midway between the highest and lowest sample, a condition which is very rarely found in practice. The results so obtained are almost always far too high, i.e., unfair to the tenant. In an acre field of rice bearing a mean yield of 15 maunds it might well be possible to find an area of 1-20th of an acre yielding 40 maunds an acre, but it would be obviously impossible to find another bearing *minus* 10 maunds an acre. The method is not far different from attempting to estimate the average income of a population by taking half the sum of the incomes of the poorest man and the richest man in it.

A second method is that at present prescribed. An officer of some standing visits the tract, of whose yield he is required to form an estimate, and selects a fairly large area, usually 1-10th of an acre, as containing an average crop. This he cuts and carries and in due course threshes and weighs. If he has time he makes two or more experiments, but it is obvious that he cannot usually manage to make a large number of such experiments. There are two main objections to this method: the first is that it depends entirely for its accuracy on the ability of the officer to select. He has to make his selection from a large number of fields growing different varieties at different points of maturity. It is very easy to be misled as to yield by the strength of straw, heavy straw being by no means always correlated with heavy yield of grain. Again it is difficult to give proper weight to the fields which will give little or no yield at all, and even in a normal year such fields are by no means rare. These factors make accurate selection very difficult. The second objection is even more important, viz., that there is no possible way of estimating what is the probability that the result of such selections is within a given range from the true mean yield. The method is comparable to estimating the average income of the population of a town by watching the streets for a few days and then picking out a man, who looked to be in average circumstances and discovering what his income is.

SAMPLING FOR RICE YIELD IN BIHAR AND ORISSA.

The only way in which a satisfactory estimate can be formed is by as close an approximation to random sampling as the circumstances permit, since that not only gets rid of the personal element of the experimenter, but also makes it possible to say what is the probability that the result of a given number of samples will be within a given range from the true mean. To put this in definite language, it should be possible to find out how many samples will be required to secure that the odds are at least 20 to 1 on the mean of the samples being within one maund of the true mean.

This is the problem to be solved. Before describing the progress made towards a solution, it is desirable to explain some terms used in modern mathematical statistics, on which the solution depends. These will, no doubt, be quite familiar to many of my readers, to whom I apologize for the digression. The first term is the arithmetical "mean", which is the real object of our search. It is, of course, the sum of all the individual results of the cuts or samples made divided by the number of cuts or samples. The true "mean" is the figure which would be obtained if the number of samples were increased, so as to cover the whole subject matter of sampling exhaustively. The whole subject matter of sampling is known as the "population", viewed as made up of a very large number of samples with varying values. The manner in which the values of the samples are numerically arranged is called the "distribution". Clearly the distribution may in some populations be very much wider than in others. The number of samples having a given value, or having a value lying within given limits, is called the "frequency" for that value or for the range of those limits. The range of limits is known as the "class interval", and, for convenience of calculation, values of samples, when these vary continuously, are lumped together in equal class intervals and treated as if they all fell at the middle point of their class interval. A most important term is the "standard deviation" of the distribution. This is a measure of the amount of dispersion or "scatter" about a particular value, usually about the mean. It is a perfectly definite value expressed in the same unit as that in which the value of each sample is expressed. It is obtained as follows. The value of each sample differs by a definite figure from the particular value, about which the dispersion is sought, *e.g.* from the mean. This difference is squared and (where the samples have been lumped together in class intervals) multiplied by the frequency of the class interval. The results are added together and divided by the sum of all the frequencies, and the square root of that result is the standard deviation. A simple illustration

SAMPLING FOR RICE YIELD IN BIHAR AND ORISSA.

will explain the process. Suppose that 8 samples give the following results.

Frequency	Value	Difference from the mean which is 6	Square of the difference multiplied by the frequency
1	4	-2	4
4	5	-1	4
1	6	0	0
0	7	+1	0
0	8	+2	0
2	9	+3	18
TOTAL 8			TOTAL 26

The standard deviation is the square root of 26 divided by 8 or 1.8 nearly.

There remains to be explained yet one more term, "the normal curve of error". This is a graphic representation of a certain class of frequency distribution, which has, among other properties, that of symmetry about the mean. To this class many of the distributions actually found in practice more or less closely conform. It has the very great advantage that the chance that a particular sample will fall within a definite range on either side of the mean can be calculated at once from standard tables, and this is what makes it so widely used in statistical practice. The properties of other classes of frequency distributions have been investigated, but their use involves difficult and laborious calculation, which makes them unsuitable for many kinds of work. It is not infrequently assumed that the determination of the standard deviation of any frequency distribution at once enables one to say what are the odds that the mean of sampling falls within given limits on either side of the true mean. This is untrue, and it is always necessary to see that the distribution under scrutiny conforms closely to the normal curve of error, before making deductions, which are true for that distribution but not for others.

Evidently then, before a practical method of obtaining a satisfactory value for the mean yield of the rice crop from a system of sampling could be recommended, it was necessary to get a clear idea of the distributions, which random sampling will give. The first attempt was made under the direction of Mr. Dobbs, Director of Agriculture, Bihar and Orissa, in 1921 in the Nawada thana of Gaya. His method was

to divide up the map into equal squares, in order to pick out the villages in which samples were to be taken, and then to sample the fields bearing the survey number 50, or the nearest fields planted with rice. For this method it was necessary to find out the date on which the crop on each field would be ready for harvest and to arrange to visit the village on that date. The practical difficulties involved made me search for another method, more automatic in the distribution of sampling in time and space. It is unfortunately impossible to use the result of Mr. Dobbs' experiment, because three samples were taken in each field and the results lumped together. This has disguised the frequency distribution so much as to make deductions extremely hazardous.

In 1923 with the help of Mr. Davies, the Settlement Officer of the Santal Parganas district, 400 random samples were obtained in a part of Godda thana. The sampler spent from the 4th December to the 22nd December travelling from village to village through the area comprising about 100 square miles. Wherever he found, on the day of his visit, the crop being actually harvested, he took a sample. In the following year the same method was applied under the supervision of the Settlement Officers, Messrs. Davies, Mansfield and Gokhale, in the Godda thana of Santal Parganas, the Jajpur thana of Cuttack, and the Purulia thana of Manbhum. The only difference was that four samples instead of one were taken, where practicable, on each field in which the harvest was in progress.

In 1925 I obtained the permission of the Bihar and Orissa Government to continue and expand the experiment in Santal Parganas and Orissa. An officer was employed in each of eight subdivisions, averaging nearly 1,000 square miles in extent, and in each some 15 centres were taken, so as to cover as widely as possible the subdivisional area. At each of these the officer was to spend four days, moving out on the first day two miles north, turning to the right and returning from the north-east, and on the succeeding days traversing the other three quadrants in the same manner. The aim was to distribute the sampling as regularly as possible both *in time* throughout the period of harvest and *in space* over the area sampled. The limitation in distance to be traversed secured that the number of samples obtained would be few in tracts where harvesting was sporadic and many in tracts where it was in full swing, thus correcting any tendency to over-emphasize the yields from relatively unimportant types. The instruction to take four samples from each field was withdrawn for reasons given below. For the supervision of this year's sampling I am indebted to Messrs. Mansfield and Houlton.

In all three years the actual process of taking each sample has been the same. The apparatus consists of a small solid block of wood, on which are fixed four pairs of guides defining two lines at an angle of 60° to one another. Along these lines through the guides are pushed two narrow wooden battens, which pass among the stalks of the paddy. Each bears a stop, so that, when this rests against the first pair of guides the length from the point where the battens cross to the slots near the ends of the battens is $67\frac{1}{4}$ inches. From one of these slots a third batten is pushed through the stalks to the slot on the other batten, and, when this third batten is fixed in the slots, an equilateral triangle is formed comprising an area of $1/3,200$ of an acre. The paddy stalks just outside the battens are bent back, and the crop inside cut and threshed out there and then. The grain is placed in a numbered bag and dried for three or four days and then weighed. The size of the triangle has been chosen to give one maund an acre for every standard tola of the sample, while the triangular shape is intended to avoid the error caused by either including or excluding a whole line of the planted paddy. The weight can be accurately determined to half a tola in ordinary country scales by double weighing. The sampler is instructed to go to the centre of the *ail* (field boundary) roughly parallel to the line on which the harvest is proceeding and take three paces into the field before laying down the solid block of wood which guides the battens. He is warned against the mistake of putting the block down so that the triangle of battens encloses it instead of excluding it, a mistake to which beginners are liable.

The results obtained in the last three years are given in Tables I and II. Besides those, to which reference has already been made, there are a set of samples taken in 1925 on the Kanke Agricultural Farm in Ranchi district and a set taken in the Atri thana of Gaya district in the same season. The system has thus been tried in the Orissa deltaic tract, on the Chota Nagpur plateau of which Ranchi forms a part, in the broken country of Santal Parganas and Manbhum, and in the alluvial Gangetic plain of North Santal Parganas and Gaya. It has been put in practice in tracts, where the harvest has been plentiful and in others, where it has been scanty. In 1925 the sampling extended in seven tracts over the whole period of the rice harvest, excluding only the very early *bhadoi* and the summer rice, reaped in March, which are harvests of very little importance. Conclusions that can be drawn from the figures may, therefore, be applied with safety to the whole province except perhaps the North-Gangetic tract.

I have relegated to the Appendix the technical discussion of the validity of my conclusions, which will be appreciated only by those readers, who have made a somewhat extensive study of statistical theory. It is enough to say here that the mean yield on the harvested area can, I think, be determined for a tract of about 1,000 square miles with an accuracy such that it will be out by more than one maund per acre in only one case out of twenty. It will be very seldom out by more than a maund and a half. To secure such determination, it is necessary to fix 12 centres, spread as evenly as possible over the area, and to put down against each centre the day on which sampling is to be done. These days should be spread evenly over the period of harvest. The sampler should go out a fixed distance in one direction and, circling round, return from another direction, so that it is secured that he covers approximately the same area on each day. He should cut one sample from each field, where he finds harvesting in progress. He will be able ordinarily to get from 30 to 40 samples in a day when the harvest is in full swing and 10 to 20 when it is slack. It is certainly not impossible to supply samplers for this work, which is extremely simple once it is explained. The men employed hitherto have been young men fresh from college. Any gazetted officer and most other officers could do it quite well. In fact, I have found that peons attached to the sampling officers fully understand what has to be done after a week or so. It is not in the least essential that the same officer should carry out all the sampling of a tract.

The method is applicable to small or large tracts. The same number of centres will probably give a slightly greater degree of accuracy for a smaller and a slightly less degree for a larger tract, because the standard deviation on the whole increases with the size of the tract, though not at all rapidly. Thus the standard deviation of the whole of the Santal Parganas is only 9.26 maunds, actually less than that of Deoghar, though greater than that of any other subdivision. To determine the mean yield of a village or an agricultural farm it would probably be advisable to make at least 10 cuts on each of 10 days. It would be better to make at least 20 cuts. (It may be noted that the standard deviation of Kanke in 1925 was nearly the highest figure found so far.)

The method could also be applied to other crops, especially cereals. It would be more difficult to apply it to crops, for which the marketable produce is obtained by processes more elaborate than threshing, or is bulky and awkward to weigh, such as jute or sugarcane. It is also an advantage if the crop is common enough to enable a fair number of samples to be taken on one day.

It is no advantage to take a large number of samples from places very close together, where the crops will naturally be very much the same on the same day. The degree of accuracy is not seriously improved by such practice. This explains why there is no need to take large samples instead of the handy samples obtained by my method. A sample of one-tenth of an acre is merely 320 of my samples taken in juxtaposition. It simply gives a determination of the mean yield of that particular field, which is not more effectively accurate than that given by say four small samples. Even four samples instead of one are not worth while, because in the great majority of cases they do not differ among themselves enough to affect the mean or the standard deviation of the whole set of samples. This may be illustrated from the columns for "all classes" and "1st cutting" for Santal Parganas, Godda thana 1924. Technically speaking, there is very high correlation between the individuals of such groups of samples, which makes the ordinary rule, that the standard deviation of the mean is the standard deviation of the population divided by the square root of the number of samples, quite inapplicable.

A feature of all the frequency distributions obtained is their tendency to have a fairly level crest, the frequency not varying greatly for some four or five class intervals around the mean. This emphasizes the difficulty of "selecting an average field", the present method of experimenting. There will be nearly as many places, where the crop is 4 maunds above or below the mean, as there will be where it is within one maund of it. Thus from the very nature of the rice crop, apart from the physical impossibility of surveying the whole of it simultaneously or nearly so, and the personal error of the experimenter, the present method of estimating is likely to be very far from accurate.

My method gives a reasonably close approximation for the mean yield of the *harvested area*. Consequently to get the gross yield it is necessary to determine what is that harvested area. It is evidently not the area planted or sown, since in practically all years some fields bear no crop worth cutting. It also excludes the field ridges (*ails*), which are usually measured in with the fields at a cadastral survey. Since the cadastral survey figure is the basis for the cropped area of the year, this second point is of importance. No very satisfactory determination of the area occupied by ridges has yet been made, but it would appear to vary between nearly 10 per cent. in hilly country to scarcely more than 2 per cent. in level country. It is usually taken as 5 per cent.

It is much more difficult to say what is the range of variation between the planted (or sown) area and the harvested area. In a bad year, when the rains stop early in September, it may well be 50 per cent. all over a subdivision. The planted area will also vary from year to year. At present we depend on estimates, which are originally those of the village chaukidar, though they are much more capable of intelligent check by higher authorities than estimates of yield are. It is possible, for instance, by driving through a tract in a motor car to get some idea of the extent to which the rice crop is going to fail altogether. It is usually believed that the estimate of cropped area is sufficiently accurate. It may be, but the belief is intuitive rather than rational.

It would certainly be a great advantage if the harvested area could be obtained by some form of sampling. A method was tried in 1925 in Santal Parganas. The results, if not very encouraging, are interesting. The plan was to make the sampler march from centre to centre across country as near as he conveniently could in a straight line. After certain intervals of time he had to count 100 paces and note how many of these ended on harvested rice land. (There is no serious difficulty in doing this in January and early February, since little rice land is reploughed by then.) Each count of 100 would give a definite percentage, in many cases 0, in many nearly 100, the difference from the full 100 being due to the presence of field ridges. The mean of these percentages would give the percentage of harvested rice land in the total area of the tract sampled. Six men were employed for nearly a month each and between them made 1971 counts. From these a mean of 47.22 per cent. was obtained. The standard deviation was 37.10. This means that it is about 21 to 1 that the true percentage is between 49.4 and 45. The accuracy obtained is hardly sufficient to justify the employment of six men for a month, in each district. Further the work is strenuous and tedious, and it is probable that it would be in practice shirked, and results fudged. The difficulty of excluding large patches of thick jungle or other inaccessible country both from the sampling and the sampled area has yet to be overcome. But it is still possible that some method on similar lines may prove practicable.

There is just one other caution which I would address to persons who have to estimate gross yields. That is that yields of rice are ultimately based, however imperfectly at present, on yields of paddy. A standard figure for the loss of weight by husking is taken, usually one-third. It is probable that this varies a good deal with the kind of paddy, and there is good reason to believe that in Orissa the loss is considerably more than one-third. It would be well if the Agricultural Department

investigated this point on their farms. If any kind of paddy husks out much below the usual, it is clearly not such a good economic proposition, as an excess of unhusked yield might suggest. Possibly the plant breeders may be able to produce a paddy, which loses less than one-third in husking, and, if so, the benefit might well be as great as that obtained from increased yield.

My general conclusions are that sampling by my method on 12 days throughout the harvest from different centres would secure a figure for the mean yield of a harvested area of 1,000 square miles correct within one maund in about 95 per cent. of cases, and that with the existing revenue and agricultural staff there would be no serious difficulty in substituting this method for the unscientific method of sampling now prescribed.

To the officers I have mentioned, under whose direction the samples were taken, and to the staff employed by them on the work I offer my cordial thanks, as I do to Mr. Dobbs for his advice and encouragement.

APPENDIX.

It is often assumed that if a number (n) of samples are taken and the standard deviation of the whole population (σ_o) is known, then the standard deviation (σ_m) of the mean of n samples is given by $\sigma_m^2 = \frac{\sigma_o^2}{n}$. This however, presupposes (1) that the samples are taken by the process of "simple" or "random" sampling, *i.e.*, that every portion of the population is equally likely to be sampled every time, and (2) that the distribution of the population is such that the distribution of the means of n samples closely approximates to the "normal" curve of error.

I have attempted to investigate both these points before recommending the method already tried for general application. The application of "simple" sampling to the rice crop would involve picking out the places for sampling by some random method such as putting numbered cards representing all the rice plots of the area in a box and drawing out the required number of cards, and then arranging to visit the plots at the time when the crop is ripe. Such a method is, I consider, impracticable. My method is practicable but is not "simple" sampling. There is some degree of selection. It has to be seen whether the adoption of it in place of "simple" sampling increases or decreases the rate at which the standard deviation of the mean diminishes as the number of cuts rises.

If an equal number of cuts were taken on each day, say 20, the process would be equivalent to (1) taking one cut from each separate tract to make up a sample, and then (2) taking the mean of 20 such samples; the "tract" being the country which the sampling officer can cover from a centre. Process (1) theoretically gives a set of samples with means having a standard deviation given by $\sigma_{m_1}^2 = \frac{\sigma_o^2}{n_1} - \frac{s_m^2}{n_1}$ where σ_o is again the standard deviation of the population, s_m that of the mean values of the tracts and n_1 the number of the tracts. Process (2) is simple sampling of the means of the samples obtained by process (1), and will give a mean with a standard deviation given by $\sigma_{m_k}^2 = \frac{\sigma_{m_1}^2}{k}$ where k is the number of samples grouped together (in this case 20).

To test the theory I have taken from the original record of Rajmahal 1925 the first eight cuts of each day, writing the 32 cuts of the first four days in one row, the 32 cuts of the second four days in the next row, and so on. (In a very few cases, where less than eight cuts were taken in a day, I have borrowed from the last cuts of the previous day or the following

day.) I have thus obtained 32 columns with 11 figures in each corresponding to the 11 centres, each surrounded by four quadrants. I have totalled the columns and taken the means. I have then taken the means of successive pairs, fours, and eights of the means so derived. I have also calculated s_m^2 by taking the means of the cuts made from each centre. I have done the same thing for Bhadrak 1925, which had 12 centres instead of 11. The results are as follows (standard deviation in maunds).

	σ_0^2	s_m^2	$\sigma_{m_1}^2$	$\sigma_{m_2}^2$	$\sigma_{m_4}^2$	$\sigma_{m_8}^2$
Rajmahal theoretical . . .	69.06	34.21	3.17	1.58	.79	.40
" observed	1.62	1.08	.40	.31
Bhadrak theoretical . . .	33.18	9.46	1.98	.99	.50	.25
" observed	1.10	.74	.54	.41

(The theoretical value of $\sigma_{m_1}^2$ is $\sigma_0^2 - s_m^2$ divided by 11 for Rajmahal and 12 for Bhadrak. The theoretical values of $\sigma_{m_2}^2$, $\sigma_{m_4}^2$ and $\sigma_{m_8}^2$ are one half, one quarter and one-eighth of $\sigma_{m_1}^2$.)

The observed results agree sufficiently well with the theory, and it seems quite clear that the standard deviation is diminished and not increased by taking samples on this plan instead of by simple sampling.

But this plan would require an equal number of cuts on each day, and this would certainly overweight the slack periods of harvest. My method avoids this, but by so doing it almost certainly increases the standard deviation of the mean as compared with that obtained by the plan just described. I cannot find any way of determining mathematically to what extent it does so, but the following test shows reasonably well what happens.

I examine below the means obtained from the cuts made to the north-east of each centre against those obtained from the cuts made to the south-east, south-west and north-west. These are given below. (Some cuts used in the general table were neglected for this purpose, because only one quadrant was sampled.)

—	Average No. of cuts in a quadrant	1st quadrant	2nd quadrant	3rd quadrant	4th quadrant	All	STANDARD DEVIATION	
							Actual	Theoretical
All S. P. . .	1432½	16.47	16.85	17.32	17.42	17.02	.25	.25
Deoghar . . .	280	20.61	19.12	21.21	20.23	20.15	.72	.69
Dumka . . .	122	19.57	20.52	19.16	20.63	20.06	.44	.41
Godda . . .	342½	12.64	14.05	15.05	15.13	14.25	.65	.35
Jamtara . . .	216	13.60	14.39	12.09	13.43	13.11	.78	.52
Rajmahal . . .	312½	21.08	21.38	22.06	22.30	21.67	.37	.47
Pakur . . .	159½	10.95	10.80	11.69	11.11	11.07	.49	.52
Bhadrak . . .	288½	13.13	14.60	13.32	13.29	13.65	.49	.34

The "failures", i.e., results falling more than one maund away from the mean of the four quadrants are underlined. The actual standard deviation is that derived from the means of the four quadrants, Sheppard's correction being applied. The theoretical standard deviation is the result of dividing the standard deviation of the whole series of cuts in the area by the square root of the average number of cuts in the quadrants. The "failures" look high at first sight, but that is due to the fact that the number of cuts in a quadrant was often distinctly lower than the standard of 20 cuts at 12 centres, which would seem to be what is required to ensure a mean correct within one maund in 95 per cent. of cases. A comparison of the actual with the theoretical standard deviation shows that on the whole my method of sampling gives about as stable a mean as simple sampling. The variation in the number of cuts at the different centres about wipes out the greater stability gained by distributing the cuts to the different centres.

It is impossible to demonstrate by mathematical processes precisely what the odds are, and, without a much wider series of tests than have yet been made, empirical deductions must be somewhat uncertain. But I think it is quite safe to say that if at least 10 and not more than 40 cuts were made from each of the 12 centres at even intervals over the harvest season, the mean yield for an area of about 1000 square miles could be determined within one maund per acre nineteen times out of twenty. The "failures" would very seldom be more than $1\frac{1}{2}$ maunds out. This depends on the assumption that the frequency distributions actually found are such that the means of a large number of cuts will form a frequency distribution closely approximating to the "normal curve of error". Where the sampling is "simple", this is certainly true of populations following the frequency distributions obtained. If, for example, 400 samples were taken by "simple" sampling from populations like the Deoghar 1925 and the Godda 1925 harvests, the spread of the mean of the 400 samples would compare with the spread of the corresponding "normal" curve (i.e., that having the same standard deviation) as follows.

Limits.	Deoghar	Normal
± .425 maunds .	1125 inside to 1000 outside	1138 inside to 1000 outside.
± .550 " .	585 " to 100 "	583 " to 100 "
± 1.275 " .	364 " to 10 "	330 " to 10 "
± 1.700 " .	342 " to 1 "	272 " to 1 "
	Godda	Normal
± .3428 " .	2517 inside to 1000 outside	2531 inside to 1000 outside.
± .5142 " .	822 " to 100 "	837 " to 100 "
± .6856 " .	3011 " to 100 "	3050 " to 100 "
± .8570 " .	1357 " to 10 "	1375 " to 10 "

It cannot then be said with certainty, what effect on the rapidity of the approach to normality would arise from the substitution of my method of sampling for "simple" sampling. But as it has little or no effect of diminishing or increasing the standard deviation of the mean, it probably has the effect of bringing the higher constants about as quickly towards their values for the normal curve. The nearest approach to a discussion of this problem is to be found in a paper by "Student" in Vol. VII pages 210 to 214 *Biometrika*. But the problem before him was not precisely mine, and he distinctly states that his problem is essentially a "small sample" problem. It deals with means of 25, not as mine with means of 250 or so. The higher constants for some of the distributions are given below (Professor Pearson's notation).

	β_1	β_2
Godda 1925	527	4.116
Deoghar 1925	1.227	4.066
Pakaur 1925	1.548	6.349
Godda 1924 1st cuttings	460	2.729
Jajpur 1924 1st cuttings	694	3.388

Professor Bowley's tables on pages 271 and 303 of his "Elements of Statistics" enable one to show how the observed mean is likely to be about the true mean. Assuming 256 samples for each tract, 10,000 trials would be distributed thus:—

(α is Professor Bowley's measure of skewness and is $\mu_3 \sigma^3$ in Professor Pearson's notation.)

—	σ	α	Not more than 1 maund above mean.	Not more than 1 maund below mean.	More than 1 maund from mean.
Godda 1925	6.63	64	4.888	4.956	156
Jamtara 1925	7.70	78	4.752	4.844	404
Pakaur 1925	6.54	86	4.900	4.982	118
Rajmahal 1925	8.31	26	4.712	4.752	536
Deoghar 1925	11.59	1.11	3.999	4.225	1,776
Dumka 1925	8.15	59	4.705	4.782	512
Jajpur 1925	9.09	1.51	4.536	4.714	750
Bhadrak 1925	5.76	69	4.943	5.003	54

It may, of course, be argued that my method, even if the same or nearly the same mean were obtained in a large number of repetitions, may be defective owing to bias. That must remain a matter of opinion.

The method does, however, secure that the cutting is well distributed in space over the area to be sampled and in time over the period of the harvest. By laying down that approximately the same area shall be under sample on each of the days of sampling, it weights the importance of the contribution made to the mean by the harvest of the day. A day in the height of the harvest will contribute 30 to 40 samples; in the slack time only 10 to 20. There is no other obvious form of bias, unless, of course, centres are selected with grave disregard to their suitability.

If it be admitted that the distribution of the means of a fairly large number of cuttings will closely approximate to a "normal curve of error" and that the standard deviation of the means is approximately that of the population divided by the square root of the number of cuttings, then the degree of accuracy can be calculated. Leaving aside the standard deviations of rice grown on particular classes of land, which are sometimes very low, it may be taken that the range of the standard deviation of rice in Bihar and Orissa is from 4 to 12 maunds. At the lower limit, 240 cuttings would give means of which only 15 out of 100,000 would be one maund or more away from the true mean. At the upper limit about 19 out of 100 would be so divergent, and about 5 out of 100 would be more than $1\frac{1}{2}$ maunds away. With a standard deviation of 8 maunds, 240 cuttings would give a mean correct within one maund about 95 times out of 100. This last is the basis of my scheme of 12 centres, from which may be expected at least 240 cuttings, but probably as many as 300. Where experience shows that the standard deviation of a tract is much higher than 8 maunds, the number of centres should be increased, if it is desired to obtain the same degree of accuracy.

It is of interest to consider what degree of accuracy would attach to a calculation of the gross yield of the whole province of Bihar and Orissa, based on the system I recommend, the harvested area being estimated as it is present. If it may be assumed that the standard deviation of the error in estimating the harvested area is about 1.20 of that estimate, it is practically certain (1,000 to 3) that the calculation of the gross yield is not more than 2 per cent. out.

TABLES

TABLE I.
Frequency distributions of rice yields, 1923-24.

Class intervals in maunds.	Santal Parganas, Godda, Jharkhand, 1924	SANTAL PARGANAS, GODDA THANA, 1924					MANERUM, PURULIA THANA, 1924					CUTTACK, JAJPUR THANA, 1924	
		1st class lands	2nd class lands	3rd class lands	All classes	1st cutting	1st class lands	2nd class lands	3rd class lands	All	1st cutting	All	1st cutting
		3	4	5	6	7	8	9	10	11	12	13	14
0-2½	55	12	1	26	39	8	..	1	2	3	1	28	16
2½-4½	24	25	24	48	100	28	1	2	5	8	2	68	37
4½-6½	44	69	49	82	200	43	6	8	40	54	18	102	45
6½-8½	45	128	53	90	271	71	17	11	74	102	33	124	43
8½-10½	71	126	86	47	259	82	18	17	84	119	87	115	42
10½-12½	27	175	78	30	283	62	18	18	100	136	42	124	62
12½-14½	18	223	50	80	342	79	17	31	78	126	39	83	34
14½-17	21	162	47	23	22	67	22	28	77	127	50	87	37
17-19½	19	173	27	4	204	57	31	19	70	120	40	52	14
19½-21½	22	118	12	4	134	49	13	27	33	73	27	39	21
21½-23½	15	51	9	3	63	17	18	16	22	56	23	24	5
23½-25½	14	16	6	3	25	4	16	11	18	40	20	28	11
25½-27½	12	4	3	..	7	4	10	8	9	27	15	21	11
27½-29½	4	6	6	..	12	3	10	5	4	19	10	17	4
29½-31½	1	2	2	..	4	1	7	8	2	17	10	6	5
31½-34	4	6	6	3	4	3	..	7	3	4	1

SAMPLING FOR RICE YIELD IN BIHAR AND ORISSA.

194

	2	1	2	1
34-36½	0	0	2	1
36½-38½	0	1	1	1
38½-40½	1
40½-42½	0
42½-44½	1
44½-46½
46½-48½
Total number of cuttings.	400	1,329	462	380
Mean in maunds	10.67	13.83	11.74	11.81
Standard deviation in maunds.	8.06	5.30	5.88	7.20
Period of cutting.	4th December to 13th December.	14th Nov. to 13th January.	16th Nov. to 27th December.	16th Nov. to 2nd January.

The "chasses" of lands are those used for rent settlement purposes, the first class being generally bottom valley lands, with a fairly secure supply of water, the third class being lands at the top of the slopes, and the second class intermediate.

SAMPLING FOR RICE YIELD IN BIHAR AND ORISSA.

21

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100	1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200	1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300	1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400	1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423	1424	1425	1426	1427	1428	1429	1430	1431	1432	1433	1434	1435	1436	1437	1438	1439	1440	1441	1442	1443	1444	1445	1446	1447	1448	1449	1450	1451	1452	1453	1454	1455	1456	1457	1458	1459	1460	1461	1462	1463	1464	1465	1466	1467	1468	1469	1470	1471	1472	1473	1474	1475	1476	1477	1478	1479	1480	1481	1482	1483	1484	148
--	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----

TABLE II—cont'd.
Frequency distribution of rice yields, 1925—cont'd.

Class intervals in mounds	Deogarh Thana				Dumka Thana				Jajpur Thana (cuttack)		Bladrak Thana	Kanke farm (Ranch)	Atri Thana (Gaya)
	1st	2nd	3rd	All	1st	2nd	3rd	All	All	All			
0-2½	0	0	2	2	0	0	0	0	31	2	0	6	
2½-4½	0	1	13	14	0	0	3	3	64	36	1	8	
4½-6½	1	4	47	52	0	1	13	14	79	68	1	15	
6½-8½	10	10	60	80	0	3	16	19	79	181	1	32	
8½-10½	16	28	58	102	0	9	18	25	79	142	2	21	
10½-12½	25	33	42	100	5	10	13	28	68	138	2	26	
12½-14½	17	37	19	73	8	18	13	44	52	186	4	14	
14½-16½	39	40	19	98	10	22	8	49	60	171	5	49	
16½-18½	32	39	21	92	30	14	20	64	41	99	6	21	
18½-20½	48	35	14	97	20	22	7	49	39	75	3	25	
20½-22½	31	22	10	63	25	15	5	45	29	35	3	23	
22½-24½	33	11	5	49	26	15	5	46	17	27	2	30	
24½-26½	34	10	7	51	18	6	6	30	7	18	6	23	
26½-28½	39	14	2	55	12	0	1	13	8	12	6	2	
28½-30½	19	5	..	24	13	0	1	14	2	5	7	14	
30½-32½	20	11	..	37	16	0	0	16	6	4	4	5	
32½-34½	27	4	..	31	10	2	2	14	4	..	0	3	

SAMPLING FOR RICE YIELD IN BIHAR AND ORISSA.

23

364—384	8	5	13	2	1	3	5	1	3	2	1	326
384—404	16	0	16	2	—	2	5	—	2	1	2	
404—424	15	1	16	3	—	3	3	—	2	2	2	
424—444	6	..	6	5	—	5	0	—	1	1	1	
444—464	9	..	9	2	—	2	2	—	2	2	1	
464—484	9	..	9	0	—	0	3	—	1	1	1	
484—51	6	..	6	1	—	1	0	—	2	2	1	
51—534	5	..	5	..	—	..	2	—	..	0	1	
534—554	6	..	6	..	—	—	
554—574	6	..	5	..	—	—	
574—594	5	..	6	..	—	—	
594—614	5	..	5	..	—	—	
614—634	1	..	1	..	—	—	
Total number of cuttings	494	310	1,123	217	138	134	637	1,152	72	326		
Mean in mannds.	27.07	18.13	20.12	24.04	18.08	14.21	12.88	13.63	26.63	17.46		
Standard deviation in mannds.	11.71	7.25	11.50	7.66	5.57	6.80	9.00	5.76	11.50	8.73		
Period of cutting	October 20th—December 26th	October 20th—December 26th	October 25th—December 21st	October 25th—December 21st	November 23rd—December 31st	October 28th—January 11th	December 4th—January 1st		

MOICU-M-18-5-6-47-27-759

